

Impurity in a Bose-Einstein condensate and the Efimov effect

Jesper Levinsen,¹ Meera M. Parish,^{1,2} and Georg M. Bruun³

¹*School of Physics and Astronomy, Monash University, Victoria 3800, Australia*

²*London Centre for Nanotechnology, Gordon Street, London, WC1H 0AH, United Kingdom*

³*Department of Physics and Astronomy, Aarhus University, DK-8000 Aarhus C, Denmark*

(Dated: October 21, 2015)

We investigate the zero-temperature properties of an impurity particle interacting with a Bose-Einstein condensate (BEC), using a variational wavefunction that includes up to two Bogoliubov excitations of the BEC. This allows one to capture three-body Efimov physics, as well as to recover the first non-trivial terms in the weak-coupling expansion. We show that the energy and quasiparticle residue of the dressed impurity (polaron) are significantly lowered by three-body correlations, even for weak interactions where there is no Efimov trimer state in a vacuum. For increasing attraction between the impurity and the BEC, we observe a smooth crossover from atom to Efimov trimer, with a superposition of states near the Efimov resonance. We furthermore demonstrate that three-body loss does not prohibit the experimental observation of these effects. Our results thus suggest a route to realizing Efimov physics in a stable quantum many-body system for the first time.

The underlying few-body dynamics in quantum many-body systems is fundamental to our understanding of nature. For instance, Fermi liquid theory is a spectacularly successful description of interacting many-body systems in terms of the dynamics of single particles, denoted quasiparticles [1, 2]. Likewise, tightly bound two-body molecules in a molecular BEC are smoothly connected to Cooper pairs in a Bardeen-Cooper-Schrieffer superfluid with decreasing attraction, as has been beautifully demonstrated in atomic gas experiments [3–5]. It is thus natural to ask whether three-body bound states lead to any observable effects in a many-body environment. In particular, Efimov famously demonstrated that there are an infinite number of three-body bound states (trimers) for identical bosons with a short range resonant interaction characterized by an infinite scattering length [6, 7]. It took more than thirty years before these Efimov states were first observed as a feature in the three-body loss rate in non-degenerate atomic gases [8–12], and more recently through Coulomb explosion imaging of ^4He trimers [13]. An intriguing question is whether Efimov states can have observable effects on a quantum many-body system, particularly outside the regime of strong three-body losses. So far, there have been no experiments reporting such effects.

Here, we show how highly imbalanced two-component atomic gases can provide a different, many-body signature of Efimov physics. The study of population imbalanced gases has already yielded fundamental new insights concerning quasiparticles in Fermi systems (Fermi polaron) [14–17]. With the recent identification of Feshbach resonances in Bose-Fermi and Bose-Bose mixtures [18–22], the study of quasiparticles in a BEC (Bose polaron) with tunable interactions is now within reach. Importantly, there is no reason to expect three-body correlations to be small for the Bose polaron, in contrast to the Fermi polaron, where these are suppressed by Fermi statistics in the two-component system [23, 24]. To have

enhanced three-body correlations in Fermi systems, one requires unequal masses [25], low dimensions [26–28], or at least three different species [29, 30].

The large majority of theoretical studies of the Bose polaron have been based on mean-field theory [31–35], or the Fröhlich model [36–42]. These approaches, however, do not include three-body Efimov correlations, and the same applies to recent variational [43] and field theoretical [44] studies. Only recently did a perturbative calculation demonstrate that three-body correlations indeed show up at third order in the impurity-boson scattering length a , leading to important effects such as a logarithmic contribution to the polaron energy [45].

In this Letter, we develop a variational ansatz for the Bose polaron that includes three-body Efimov physics, and in addition recovers all contributions to the energy in the weak-coupling limit up to and including the logarithmic term at order a^3 . We show that Efimov correlations significantly lower the energy and quasiparticle residue, even for scattering lengths substantially smaller in magnitude than $a_- < 0$, the scattering length where the deepest Efimov trimer crosses the three-body continuum in a vacuum. As a approaches a_- , the energy exhibits a clear avoided crossing, reflecting a smooth change in the quasiparticle character, from single-particle to three-body. We furthermore demonstrate that these effects can be observed before encountering resonantly enhanced three-body loss, which has so far been the main experimental signature of Efimov physics.

Model.— We consider an impurity particle immersed in a weakly interacting BEC with $n_0 a_B^3 \ll 1$, where n_0 denotes the condensate density and a_B the boson-boson scattering length. The addition of an impurity atom to the condensate introduces two additional length scales: The impurity-boson scattering length a , and the so-called three-body parameter. The latter sets the scale of one Efimov trimer in the universal limit of $1/a \rightarrow 0$, from which all other trimer energies can be determined. In the

cold atoms context, it is natural to use a_- as the three-body parameter. It has recently emerged in studies of identical bosons [46–48] that this parameter is universally related to the van der Waals range. Thus, we will fix a_- through the two-body physics.

For this purpose, it is convenient to use the two-channel Hamiltonian (setting $\hbar$ and the volume to 1):

$$\hat{H} = \sum_{\mathbf{k}} \left[E_{\mathbf{k}} \beta_{\mathbf{k}}^\dagger \beta_{\mathbf{k}} + \epsilon_{\mathbf{k}} c_{\mathbf{k}}^\dagger c_{\mathbf{k}} + (\epsilon_{\mathbf{k}}^d + \nu_0) d_{\mathbf{k}}^\dagger d_{\mathbf{k}} \right] + g\sqrt{n_0} \sum_{\mathbf{k}} \left(d_{\mathbf{k}}^\dagger c_{\mathbf{k}} + h.c. \right) + g \sum_{\mathbf{q}, \mathbf{k}} \left(d_{\mathbf{q}}^\dagger c_{\mathbf{q}-\mathbf{k}} b_{\mathbf{k}} + h.c. \right), \quad (1)$$

which measures the energy with respect to that of the BEC. Here, $b_{\mathbf{k}}^\dagger$ and $c_{\mathbf{k}}^\dagger$ create a boson and the impurity, respectively, with momentum $\mathbf{k}$ and single-particle energy $\epsilon_{\mathbf{k}} = k^2/2m$. The impurity interacts with a boson by forming a closed channel molecule, created by $d_{\mathbf{k}}^\dagger$, with dispersion $\epsilon_{\mathbf{k}}^d = \epsilon_{\mathbf{k}}/2$ and bare detuning ν_0 . The coupling strength for this process, g , is taken constant up to a momentum cut-off Λ . We follow the usual weak-coupling prescription, relating the boson operator b to the Bogoliubov excitation β via $b_{\mathbf{k}} = u_{\mathbf{k}} \beta_{\mathbf{k}} - v_{\mathbf{k}} \beta_{-\mathbf{k}}^\dagger$ with real and positive coherence factors $u_{\mathbf{k}}^2 = [1 + (\epsilon_{\mathbf{k}} + \mu)/E_{\mathbf{k}}]/2$ and $v_{\mathbf{k}}^2 = [-1 + (\epsilon_{\mathbf{k}} + \mu)/E_{\mathbf{k}}]/2$, dispersion $E_{\mathbf{k}} = \sqrt{\epsilon_{\mathbf{k}}(\epsilon_{\mathbf{k}} + 2\mu)}$, and chemical potential $\mu = 4\pi a_B n_0/m$. The impurity is assumed to have the same mass m as the bosons, as we do not expect our results to change qualitatively in the mass imbalanced system.

By considering impurity-boson scattering, we relate the bare molecule detuning ν_0 , coupling g , and momentum cut-off to the scattering length a and the effective range r_0 [49, 50]. The additional length scale r_0 provided by the closed channel fixes the three-body parameter of Efimov physics. In particular, by solving the impurity-boson-boson problem in vacuum with $a_B = 0$, we find $a_- \approx 2467r_0$, which is consistent with calculations involving realistic interatomic potentials for a range of heteronuclear systems [51]. This large separation of scales means we can always take $|a/r_0| \gg 1$, where the effect of the closed channel on two-body processes is negligible.

Variational approach.— We consider the wavefunction:

$$|\psi\rangle = \left(\alpha_0 c_0^\dagger + \sum_{\mathbf{k}} \alpha_{\mathbf{k}} c_{-\mathbf{k}}^\dagger \beta_{\mathbf{k}}^\dagger + \frac{1}{2} \sum_{\mathbf{k}_1 \mathbf{k}_2} \alpha_{\mathbf{k}_1 \mathbf{k}_2} c_{-\mathbf{k}_1 - \mathbf{k}_2}^\dagger \beta_{\mathbf{k}_1}^\dagger \beta_{\mathbf{k}_2}^\dagger + \gamma_0 d_0^\dagger + \sum_{\mathbf{k}} \gamma_{\mathbf{k}} d_{-\mathbf{k}}^\dagger \beta_{\mathbf{k}}^\dagger \right) |\Phi\rangle, \quad (2)$$

with $|\Phi\rangle$ the wavefunction of the interacting BEC. The first line of Eq. (2) describes the bare impurity and its dressing by one or two Bogoliubov modes. The second line describes the impurity bound to a boson and forming a closed channel molecule, which can be dressed by a Bogoliubov mode. The variational approach consists in taking the stationary condition $\langle \partial\psi | (\hat{H} - E) | \psi \rangle = 0$

where the derivative is with respect to any of the expansion parameters. This procedure yields five coupled equations for the energy E [50]. Defining

$$\eta = \frac{g^2 \sum_{\mathbf{k}} u_{\mathbf{k}} \alpha_{\mathbf{k}}}{E - \nu_0}, \quad \xi_{\mathbf{k}} = \frac{g^2 \sum_{\mathbf{k}'} u_{\mathbf{k}'} \alpha_{\mathbf{k} \mathbf{k}'}}{E - \nu_0 - \epsilon_{\mathbf{k}}^d - E_{\mathbf{k}}}, \quad (3)$$

which remain finite in the limit $\nu_0 \rightarrow \infty$, and carrying out the renormalization procedure, we finally arrive at the coupled integral equations

$$\begin{aligned} \mathcal{T}^{-1}(\mathbf{0}, E) \eta &= \frac{n_0}{E} \eta + \sqrt{n_0} \sum_{\mathbf{k}} \left(\frac{u_{\mathbf{k}} \xi_{\mathbf{k}}}{E - \epsilon_{\mathbf{k}} - E_{\mathbf{k}}} - \frac{v_{\mathbf{k}} \xi_{\mathbf{k}}}{E} \right), \\ \mathcal{T}^{-1}(\mathbf{k}_1, E - E_{\mathbf{k}_1}) \xi_{\mathbf{k}_1} &= \sqrt{n_0} \left(\frac{u_{\mathbf{k}_1} \eta}{E - \epsilon_{\mathbf{k}_1} - E_{\mathbf{k}_1}} - \frac{v_{\mathbf{k}_1} \eta}{E} \right) \\ &+ \frac{n_0 \xi_{\mathbf{k}_1}}{E - \epsilon_{\mathbf{k}_1} - E_{\mathbf{k}_1}} + \sum_{\mathbf{k}_2} \left(\frac{u_{\mathbf{k}_1} u_{\mathbf{k}_2} \xi_{\mathbf{k}_2}}{E_{\mathbf{k}_1 \mathbf{k}_2}} + \frac{v_{\mathbf{k}_1} v_{\mathbf{k}_2} \xi_{\mathbf{k}_2}}{E} \right), \quad (4) \end{aligned}$$

where $E_{\mathbf{k}_1 \mathbf{k}_2} \equiv E - E_{\mathbf{k}_1} - E_{\mathbf{k}_2} - \epsilon_{\mathbf{k}_1 + \mathbf{k}_2}$. Here, the medium $\mathcal{T}$ matrix is

$$\mathcal{T}^{-1}(\mathbf{k}_1, E) = \frac{m_r}{2\pi a} - \frac{m_r^2 r_0}{2\pi} (E - \epsilon_{\mathbf{k}_1}^d) - \Pi(\mathbf{k}_1, E), \quad (5)$$

with $\Pi(\mathbf{k}_1, E) = \sum_{\mathbf{k}_2} [u_{\mathbf{k}_2}^2 / (E - E_{\mathbf{k}_2} - \epsilon_{\mathbf{k}_1 + \mathbf{k}_2}) + 2m_r/k_2^2]$ the pair propagator in the BEC and $m_r = m/2$ the reduced mass.

A few comments on our variational approach are in order. First, had the two equations (4) not been coupled, the first of these would simply reproduce the mean-field polaron energy, $E = 4\pi n_0 a/m$ (for a non-zero r_0 there is a negligible correction $\sim \mathcal{O}(n^2 a^3 r_0)$ as r_0 is by far the smallest length scale). Second, in the limit $n_0 \rightarrow 0$ the second equation exactly yields the Efimov spectrum for the impurity and two identical, non-interacting bosons. This explicitly demonstrates that our variational ansatz contains the relevant three-body Efimov correlations. For a discussion of the three-body problem with an effective range, we refer the reader to Refs. [52–54].

Furthermore, since our wavefunction includes all two-body excitations, it recovers all second order perturbation terms in a . This is because all relevant Feynman diagrams contain at most two concurrent boson propagators (neglecting $n_0 a_B^3$ corrections) [45]. Only a subset of these diagrams are produced if one restricts the approach to one Bogoliubov mode, as in Ref. [43]. Although our ansatz does not reproduce all third order self energy diagrams, it does indeed capture the logarithmic term in the energy, since this originates from diagrams containing at most two simultaneous Bogoliubov excitations [45].

Finally, we remark that had we used a single-channel model from the outset, we would have arrived at equations identical to Eqs. (4) with $r_0 = 0$ in Eq. (5). However, in this case we would need to set the three-body parameter in some other way, for instance through the use of a Gaussian cut-off.




Figure 1. (a) Spectrum and (b) residue of the Bose polaron for $n_0^{1/3}a_- = -1$ and $a_B/a_- = -1/50$. We show the results from our variational calculation (purple, solid), and the variational wavefunction excluding three-body correlations and effective range effects (blue, dashed). The gray lines are the results of perturbation theory: Mean-field (dashed), second order [38] (dot-dashed), and third order [45] (dotted). We also display the energy of the deepest Efimov state in a vacuum (red, dashed), and for $a > 0$ the threshold of the atom-dimer continuum given by the impurity-boson bound state energy $-(\sqrt{1 - 2r_0/a} - 1)^2/(2m_r r_0^2)$ (black, solid). The insets illustrate the crossover in the wavefunction (the atoms are not drawn to scale).

Results.— The Bose polaron is characterized by its quasiparticle properties, i.e. the energy E and residue $Z \equiv |\alpha_0|^2$ [50]. In Fig. 1, we display these quantities calculated from Eq. (4) as a function of $1/(n_0^{1/3}a)$ for realistic experimental parameters. For weak interaction, $1/(n_0^{1/3}a) \ll -1$, the polaron energy is close to the mean-field value $4\pi an_0/m$ and $Z \simeq 1$. With increasing coupling, E starts to deviate from the mean-field result, whereas there is better agreement with second [38] and third [45] order perturbation theory. This is as expected since, as argued above, our variational approach contains all diagrams contributing to the energy at these orders of perturbation theory. While $Z = 1$ within mean-field theory, and the second order perturbative expression [45] is negative in the range plotted, the agreement at third order with our variational ansatz is very good, leading us to conclude that although the variational ansatz does not contain all third order terms for the residue, it likely includes the most important ones.




Figure 2. Residue of the Bose polaron as a function of $1/(n_0^{1/3}a)$ and $1/(n_0^{1/3}|a_-|)$ for $a_B/a_- = -1/50$. The white dashed line is $a = a_-$. For sufficiently small $1/(n_0^{1/3}|a_-|)$, the residue remains close to 1 around unitarity, implying that the Efimov trimer has disappeared.

Closer to resonance, three-body correlations become important, leading to an avoided crossing between a state best described as a single particle interacting attractively with the medium and an Efimov trimer dressed by interactions, as seen in Fig. 1 and illustrated in the insets. The residue correspondingly decreases as the wavefunction resembles the three-body Efimov state. This crossover is of course completely missed by the perturbative expressions. It is also missed by the variational ansatz used in Ref. [43], which includes only two-body correlations, i.e., only the first two terms in Eq. (2). Within this approach the quasiparticle residue only goes to zero once the polaron approaches the atom-dimer continuum. Our analysis is the first to include Efimov physics for the Bose polaron, which we see strongly changes its properties at a qualitative level. Even outside the regime of Efimov physics in vacuum, there are strong effects of Efimov physics: For instance, at $a = a_-/2$ the magnitude of the energy is increased by 50% when comparing with the variational approach restricted to two-body correlations.

To investigate the strength of the coupling between the two branches involved in the avoided crossing, we show in Fig. 2 the residue as a function of $1/(n_0^{1/3}a)$ and $1/(n_0^{1/3}|a_-|)$. Focussing first on the region where $1/(n_0^{1/3}|a_-|) \geq 1$, we see that the crossover region is located around $a \sim a_-$, where Z decreases to almost zero. Furthermore, the range of scattering lengths over which this transition occurs decreases with decreasing $|a_-|$, reflecting how the coupling to the Efimov state becomes weaker. This can be understood from a simple dimensional analysis as follows: Treating momentarily the Efimov trimer as a point particle, the coupling of this state to two bosons and the impurity involves four particles,

three in-going and one out-going or vice versa. The coupling strength for this process must therefore scale as Length/Mass. In the avoided crossing region, there is only one relevant length scale, $a \sim a_-$, and consequently the coupling strength $\sim a_-/m$. On the other hand, in the regime $1/(n_0^{1/3}|a_-|) \ll 1$ the perturbative energy far exceeds that of the Efimov trimer in vacuum when $a \sim a_-$ and thus the quasiparticle remains well-defined even beyond this point, as seen in the figure. Eventually, of course, the residue is suppressed as the polaron becomes dimer-like for sufficiently small and positive a .

To investigate the dependence on a_B , we plot in Fig. 3(a-b) the energy and residue of the Bose polaron for $n_0^{1/3}a_- = -1/4$ and three different values of the boson-boson scattering length. In both quantities we clearly observe how the crossover becomes sharper with increasing $a_B/|a_-|$, since boson-boson repulsion suppresses the formation of the Efimov trimer. An even larger suppression is expected to occur for any four-body Efimov state [55] consisting of the impurity and three bosons, if such states exist (these have so far only been investigated for unequal masses [56]). Indeed, the impact of four-body states on the many-body system will be further reduced since one expects them to appear for $|a| < |a_-|$, resulting in a smaller coupling to the bound state similar to that shown in Fig. 2. It is thus likely that such states can be ignored even in a regime where the coupling to the three-body state is strong.

While the energy and residue of the Fermi polaron have been measured using radio-frequency (RF) spectroscopy [14–16], the presence of the Efimov trimer in a cold atoms context has hitherto only been observed indirectly, as either a significant increase (suppression) in the three-body loss rate for negative (positive) scattering lengths [8–11], or via enhanced loss from the RF association of trimers close to the atom-dimer threshold [12]. An important question is therefore whether the shifts in the polaron energy and residue due to Efimov correlations can be observed before encountering prohibitively fast losses. To investigate this, we note that the main loss process – three-body recombination to a deeply bound molecule and an atom – is local, since it requires three atoms to be separated within a distance comparable to the size of the outgoing states. Hence, the rate of this loss is captured, within a perturbative approach, by the expectation value [57, 58]

$$\Gamma = \Delta \langle \psi | \sum_{\mathbf{k}_1 \mathbf{k}_2 \mathbf{p}} d_{\mathbf{p}-\mathbf{k}_1}^\dagger b_{\mathbf{k}_1}^\dagger b_{\mathbf{k}_2} d_{\mathbf{p}-\mathbf{k}_2} | \psi \rangle. \quad (6)$$

Making the replacement $\beta_k \rightarrow b_k$ at short range, we get $\Gamma \approx \Delta \sum_k |\gamma_k|^2$. Note that the constant Δ should be fixed by comparison with experimental data outside the regime of resonantly enhanced losses; thus, in Fig. 3(c) we plot Γ/Δ . We see that the loss rate increases with decreasing Z due to the onset of Efimov correlations, and




Figure 3. (a) Energy, (b) residue, and (c) three-body loss rate for $n_0^{1/3}|a_-| = 0.25$ and different values of $a_B/|a_-|$: 0.01 (purple, solid), 0.1 (purple, dashed), 0.5 (purple, dotted). In (a) we also display the energy of the deepest Efimov state in vacuum (red, dashed). The blue lines are the variational results excluding three-body correlations as well as effective range effects.

saturates to that of the vacuum Efimov state. Figure 3, however, demonstrates a crucial point: The reduction in the energy and residue due to three-body correlations is still substantial away from $a = a_-$, where the decay rate can be more than *two orders of magnitude* smaller than that of the vacuum Efimov state. From this we conclude that the lifetime of the impurity is indeed long enough in this regime to allow a detection of three-body effects. This mirrors previous work on trimers in a quasi-two-dimensional geometry, which argues that Efimov physics can be probed by coupling an Efimov trimer to a trimer extended within the plane [59].

Conclusions. – Using a variational ansatz which includes three-body Efimov correlations as well as the lowest non-trivial terms in the weak-coupling expansion, we demonstrated that Efimov physics dramatically changes the properties of the Bose polaron. The energy and quasiparticle residue are lowered significantly, even when there is no Efimov trimer state in a vacuum. As the critical scattering length for the emergence of the trimer state is approached, the energy exhibits an avoided crossing and the residue quickly decreases, reflecting the fact that the impurity smoothly becomes bound in an Efimov trimer. Crucially, we showed that three-body loss associated with the Efimov state does not prohibit the observation of

these effects. In conclusion, our results demonstrate how Efimov physics qualitatively changes the properties of a quantum many-body system, and how it can be observed in signals other than loss.

We gratefully acknowledge fruitful discussion with Jan Arlt, Rasmus S. Christensen, Yusuke Nishida, and Richard Schmidt. G.M.B. wishes to acknowledge the support of the Villum Foundation via grant VKR023163.

-
- [1] L. Landau, *Theory of Fermi-liquids*, Sov. Phys. JETP **3**, 920 (1957).
 - [2] G. Baym and C. Pethick, *Landau Fermi-Liquid Theory: Concepts and Applications* (Wiley-VCH, 1991).
 - [3] A. Leggett, *Diatomic molecules and cooper pairs*, in *Modern Trends in the Theory of Condensed Matter*, Lecture Notes in Physics, Vol. 115, edited by A. Pólkalski and J. Przystawa (Springer Berlin Heidelberg, 1980) pp. 13–27.
 - [4] P. Nozières and S. Schmitt-Rink, *Bose condensation in an attractive fermion gas: From weak to strong coupling superconductivity*, Journal of Low Temperature Physics **59**, 195 (1985).
 - [5] M. Randeria and E. Taylor, *Crossover from Bardeen-Cooper-Schrieffer to Bose-Einstein Condensation and the Unitary Fermi Gas*, Annual Review of Condensed Matter Physics **5**, 209 (2014).
 - [6] V. Efimov, *Energy levels arising from resonant two-body forces in a three-body system*, Phys. Lett. B **33**, 563 (1970).
 - [7] E. Braaten and H.-W. Hammer, *Universality in few-body systems with large scattering length*, Physics Reports **428**, 259 (2006).
 - [8] T. Kraemer, M. Mark, P. Waldburger, J. G. Danzl, C. Chin, B. Engeser, A. D. Lange, K. Pilch, A. Jaakkola, H.-C. Nägerl, and R. Grimm, *Evidence for Efimov quantum states in an ultracold gas of caesium atoms*, Nature **440**, 315 (2006).
 - [9] M. Zaccanti, B. Deissler, C. D’Errico, M. Fattori, M. Jona-Lasinio, S. Muller, G. Roati, M. Inguscio, and G. Modugno, *Observation of an Efimov spectrum in an atomic system*, Nature Phys. **5**, 586 (2009).
 - [10] N. Gross, Z. Shotan, S. Kokkelmans, and L. Khaykovich, *Observation of Universality in Ultracold ^7Li Three-Body Recombination*, Phys. Rev. Lett. **103**, 163202 (2009).
 - [11] S. E. Pollack, D. Dries, and R. G. Hulet, *Universality in Three- and Four-Body Bound States of Ultracold Atoms*, Science **326**, 1683 (2009).
 - [12] T. Lompe, T. B. Ottenstein, F. Serwane, A. N. Wenz, G. Zürn, and S. Jochim, *Radio-Frequency Association of Efimov Trimers*, Science **330**, 940 (2010).
 - [13] M. Kunitski, S. Zeller, J. Voigtsberger, A. Kalinin, L. P. H. Schmidt, M. Schöffler, A. Czasch, W. Schöllkopf, R. E. Grisenti, T. Jahnke, D. Blume, and R. Dörner, *Observation of the Efimov state of the helium trimer*, Science **348**, 551 (2015).
 - [14] A. Schirotzek, C.-H. Wu, A. Sommer, and M. W. Zwierlein, *Observation of Fermi Polarons in a Tunable Fermi Liquid of Ultracold Atoms*, Phys. Rev. Lett. **102**, 230402 (2009).
 - [15] C. Kohstall, M. Zaccanti, M. Jag, A. Trenkwalder, P. Massignan, G. M. Bruun, F. Schreck, and R. Grimm, *Metastability and coherence of repulsive polarons in a strongly interacting Fermi mixture*, Nature (London) **485**, 615 (2012).
 - [16] M. Koschorreck, D. Pertot, E. Vogt, B. Fröhlich, M. Feld, and M. Köhl, *Attractive and repulsive Fermi polarons in two dimensions*, Nature (London) **485**, 619 (2012).
 - [17] P. Massignan, M. Zaccanti, and G. M. Bruun, *Polarons, dressed molecules and itinerant ferromagnetism in ultracold Fermi gases*, Reports on Progress in Physics **77**, 034401 (2014).
 - [18] C.-H. Wu, J. W. Park, P. Ahmadi, S. Will, and M. W. Zwierlein, *Ultracold Fermionic Feshbach Molecules of $^{23}\text{Na}^{40}\text{K}$* , Phys. Rev. Lett. **109**, 085301 (2012).
 - [19] M.-S. Heo, T. T. Wang, C. A. Christensen, T. M. Rvachov, D. A. Cotta, J.-H. Choi, Y.-R. Lee, and W. Ketterle, *Formation of ultracold fermionic NaLi Feshbach molecules*, Phys. Rev. A **86**, 021602 (2012).
 - [20] T. D. Cumby, R. A. Shewmon, M.-G. Hu, J. D. Perreault, and D. S. Jin, *Feshbach-molecule formation in a Bose-Fermi mixture*, Phys. Rev. A **87**, 012703 (2013).
 - [21] G. Roati, M. Zaccanti, C. D’Errico, J. Catani, M. Modugno, A. Simoni, M. Inguscio, and G. Modugno, *^{39}K Bose-Einstein Condensate with Tunable Interactions*, Phys. Rev. Lett. **99**, 010403 (2007).
 - [22] K. Pilch, A. D. Lange, A. Prantner, G. Kerner, F. Ferlaino, H.-C. Nägerl, and R. Grimm, *Observation of inter-species Feshbach resonances in an ultracold Rb-Cs mixture*, Phys. Rev. A **79**, 042718 (2009).
 - [23] R. Combescot and S. Giraud, *Normal State of Highly Polarized Fermi Gases: Full Many-Body Treatment*, Phys. Rev. Lett. **101**, 050404 (2008).
 - [24] J. Vlietinck, J. Ryckebusch, and K. Van Houcke, *Quasi-particle properties of an impurity in a Fermi gas*, Phys. Rev. B **87**, 115133 (2013).
 - [25] C. J. M. Mathy, M. M. Parish, and D. A. Huse, *Trimers, molecules and polarons in imbalanced atomic Fermi gases*, Phys. Rev. Lett. **106**, 166404 (2011).
 - [26] G. Orso, E. Burovski, and T. Jolicoeur, *Luttinger Liquid of Trimers in Fermi Gases with Unequal Masses*, Phys. Rev. Lett. **104**, 065301 (2010).
 - [27] M. M. Parish, *Polaron-molecule transitions in a two-dimensional Fermi gas*, Phys. Rev. A **83**, 051603 (2011).
 - [28] M. M. Parish and J. Levinsen, *Highly polarized Fermi gases in two dimensions*, Phys. Rev. A **87**, 033616 (2013).
 - [29] Y. Nishida, *Polaronic Atom-Trimer Continuity in Three-Component Fermi Gases*, Phys. Rev. Lett. **114**, 115302 (2015).
 - [30] W. Yi and X. Cui, *Polarons in ultracold Fermi superfluids*, Phys. Rev. A **92**, 013620 (2015).
 - [31] G. E. Astrakharchik and L. P. Pitaevskii, *Motion of a heavy impurity through a Bose-Einstein condensate*, Phys. Rev. A **70**, 013608 (2004).
 - [32] F. M. Cucchiatti and E. Timmermans, *Strong-Coupling Polarons in Dilute Gas Bose-Einstein Condensates*, Phys. Rev. Lett. **96**, 210401 (2006).
 - [33] R. M. Kalas and D. Blume, *Interaction-induced localization of an impurity in a trapped Bose-Einstein condensate*, Phys. Rev. A **73**, 043608 (2006).
 - [34] M. Bruderer, W. Bao, and D. Jaksch, *Self-trapping of impurities in Bose-Einstein condensates: Strong attractive and repulsive coupling*, EPL (Europhysics Letters) **82**, 30004 (2008).

- [35] A. G. Volosniev, H.-W. Hammer, and N. T. Zinner, *Real-time dynamics of an impurity in an ideal Bose gas in a trap*, Phys. Rev. A **92**, 023623 (2015).
- [36] B.-B. Huang and S.-L. Wan, *Polaron in Bose-Einstein-Condensation System*, Chinese Physics Letters **26**, 080302 (2009).
- [37] J. Tempere, W. Casteels, M. K. Oberthaler, S. Knoop, E. Timmermans, and J. T. Devreese, *Feynman path-integral treatment of the BEC-impurity polaron*, Phys. Rev. B **80**, 184504 (2009).
- [38] W. Casteels and M. Wouters, *Polaron formation in the vicinity of a narrow Feshbach resonance*, Phys. Rev. A **90**, 043602 (2014).
- [39] A. Shashi, F. Grusdt, D. A. Abanin, and E. Demler, *Radio-frequency spectroscopy of polarons in ultracold Bose gases*, Phys. Rev. A **89**, 053617 (2014).
- [40] F. Grusdt, Y. E. Shchadilova, A. N. Rubtsov, and E. Demler, *Renormalization group approach to the Fröhlich polaron model: application to impurity-BEC problem*, Sci. Rep. **5**, 12124 (2015).
- [41] Y. E. Shchadilova, F. Grusdt, A. N. Rubtsov, and E. Demler, *Polaronic mass renormalization of impurities in BEC: correlated Gaussian wavefunction approach*, ArXiv e-prints (2014), arXiv:1410.5691 [cond-mat.quant-gas].
- [42] J. Vlietinck, W. Casteels, K. V. Houcke, J. Tempere, J. Ryckebusch, and J. T. Devreese, *Diagrammatic Monte Carlo study of the acoustic and the Bose-Einstein condensate polaron*, New Journal of Physics **17**, 033023 (2015).
- [43] W. Li and S. Das Sarma, *Variational study of polarons in Bose-Einstein condensates*, Phys. Rev. A **90**, 013618 (2014).
- [44] S. P. Rath and R. Schmidt, *Field-theoretical study of the Bose polaron*, Phys. Rev. A **88**, 053632 (2013).
- [45] R. Sogaard Christensen, J. Levinsen, and G. M. Bruun, *Quasiparticle properties of a mobile impurity in a Bose-Einstein condensate*, Phys. Rev. Lett. **115**, 160401 (2015).
- [46] M. Berninger, A. Zenesini, B. Huang, W. Harm, H.-C. Nägerl, F. Ferlaino, R. Grimm, P. S. Julienne, and J. M. Hutson, *Universality of the Three-Body Parameter for Efimov States in Ultracold Cesium*, Phys. Rev. Lett. **107**, 120401 (2011).
- [47] S. Roy, M. Landini, A. Trenkwalder, G. Semeghini, G. Spagnolli, A. Simoni, M. Fattori, M. Inguscio, and G. Modugno, *Test of the Universality of the Three-Body Efimov Parameter at Narrow Feshbach Resonances*, Phys. Rev. Lett. **111**, 053202 (2013).
- [48] J. Wang, J. P. D’Incao, B. D. Esry, and C. H. Greene, *Origin of the Three-Body Parameter Universality in Efimov Physics*, Phys. Rev. Lett. **108**, 263001 (2012).
- [49] G. M. Bruun and C. J. Pethick, *Effective Theory of Feshbach Resonances and Many-Body Properties of Fermi Gases*, Phys. Rev. Lett. **92**, 140404 (2004).
- [50] See Supplemental Material for the impurity-boson scattering analysis, variational equations, residue, and decay rate.
- [51] Y. Wang, J. Wang, J. P. D’Incao, and C. H. Greene, *Universal Three-Body Parameter in Heteronuclear Atomic Systems*, Phys. Rev. Lett. **109**, 243201 (2012).
- [52] D. S. Petrov, *Three-boson problem near a narrow Feshbach resonance*, Phys. Rev. Lett. **93**, 143201 (2004).
- [53] J. Levinsen, T. G. Tiecke, J. T. M. Walraven, and D. S. Petrov, *Atom-dimer scattering and long-lived trimers in fermionic mixtures*, Phys. Rev. Lett. **103**, 153202 (2009).
- [54] D. S. Petrov, *The few-atom problem*, in *Proceedings of the Les Houches Summer Schools, Session 94*, edited by C. Salomon, G. V. Shlyapnikov, and L. F. Cugliandolo (Oxford University Press, Oxford, England, 2013) p. 109, preprint available at arXiv:1206.5752.
- [55] J. von Stecher, J. P. D’Incao, and C. H. Greene, *Four-body legacy of the Efimov effect*, Nature Physics **5**, 417 (2009).
- [56] D. Blume and Y. Yan, *Generalized Efimov Scenario for Heavy-Light Mixtures*, Phys. Rev. Lett. **113**, 213201 (2014).
- [57] E. Braaten and H.-W. Hammer, *Three-Body Recombination into Deep Bound States in a Bose Gas with Large Scattering Length*, Phys. Rev. Lett. **87**, 160407 (2001).
- [58] J. Levinsen and D. Petrov, *Atom-dimer and dimer-dimer scattering in fermionic mixtures near a narrow Feshbach resonance*, The European Physical Journal D **65**, 67 (2011).
- [59] J. Levinsen, P. Massignan, and M. M. Parish, *Efimov Trimers under Strong Confinement*, Phys. Rev. X **4**, 031020 (2014).

Supplemental Material: Impurity in a Bose-Einstein condensate and the Efimov effect

Jesper Levinsen,¹ Meera M. Parish,^{1,2} and Georg M. Bruun³

¹*School of Physics and Astronomy, Monash University, Victoria 3800, Australia*

²*London Centre for Nanotechnology, Gordon Street, London, WC1H 0AH, United Kingdom*

³*Department of Physics and Astronomy, Aarhus University, DK-8000 Aarhus C, Denmark*

IMPURITY-BOSON SCATTERING

The bare molecule energy ν_0 is not a physical observable since the molecule is dressed by open channel impurity-boson pairs. Likewise, the coupling is assumed to be a constant g for momenta below an ultraviolet cut-off Λ , which is not related directly to any observable. We can relate these parameters to observables by considering impurity-boson scattering via the closed channel molecule. In a vacuum, the scattering matrix is [49, 58]

$$\mathcal{T}_{\text{vac}}(\omega) = \frac{g^2}{\omega - \nu_0 - g^2 \Pi_{\text{vac}}(0) - g^2 \Delta \Pi_{\text{vac}}(\omega)} \quad (\text{S1})$$

where $\Pi_{\text{vac}}(\omega) = \sum_{\mathbf{k}} (\omega - k^2/2m_r)^{-1}$ is the vacuum pair propagator with energy ω , $m_r = m/2$ is the reduced mass, and $\Delta \Pi_{\text{vac}}(\omega) = \Pi_{\text{vac}}(\omega) - \Pi_{\text{vac}}(0)$. Equating (S1) evaluated at the scattering energy $\omega = p^2/2m_r$ with the effective range expansion $\mathcal{T}_{\text{vac}}(p^2/2m_r) = 2\pi a/m_r(1 - k^2 a r_0/2 + ika)$, where r_0 is the effective range, we obtain $a = m_r g^2 (g^2 m_r \Lambda / \pi^2 - \nu_0) / 2\pi$ and $r_0 = -2\pi / m_r^2 g^2$. In this way we replace the bare parameters ν_0 and Λ with the low energy observables a and r_0 for the impurity-boson scattering. Then we are at liberty to take the limits $\nu_0 \rightarrow \infty$, $\Lambda \rightarrow \infty$.

VARIATIONAL EQUATIONS

Taking the derivative $\langle \partial \psi | (\hat{H} - E) | \psi \rangle = 0$ with respect to α_0 , $\alpha_{\mathbf{k}}$, $\alpha_{\mathbf{k}_1 \mathbf{k}_2}$, γ_0 , and $\gamma_{\mathbf{k}}$ yields the five equations

$$\begin{aligned} E\alpha_0 &= g\sqrt{n_0}\gamma_0 - g \sum_{\mathbf{k}} v_{\mathbf{k}}\gamma_{\mathbf{k}}, \\ (E - \epsilon_{\mathbf{k}} - E_{\mathbf{k}}) \alpha_{\mathbf{k}} &= g u_{\mathbf{k}}\gamma_0 + g\sqrt{n_0}\gamma_{\mathbf{k}}, \\ E_{\mathbf{k}_1 \mathbf{k}_2} \alpha_{\mathbf{k}_1 \mathbf{k}_2} &= g(\gamma_{\mathbf{k}_1} u_{\mathbf{k}_2} + \gamma_{\mathbf{k}_2} u_{\mathbf{k}_1}), \\ (E - \nu_0)\gamma_0 &= g\sqrt{n_0}\alpha_0 + g \sum_{\mathbf{k}} u_{\mathbf{k}}\alpha_{\mathbf{k}}, \\ (E - \epsilon_{\mathbf{k}}^{\text{d}} - \nu_0 - E_{\mathbf{k}}) \gamma_{\mathbf{k}} &= g\sqrt{n_0}\alpha_{\mathbf{k}} - g v_{\mathbf{k}}\alpha_0 + g \sum_{\mathbf{k}'} u_{\mathbf{k}'} \alpha_{\mathbf{k}\mathbf{k}'}, \end{aligned} \quad (\text{S2})$$

where $E_{\mathbf{k}_1 \mathbf{k}_2} \equiv E - E_{\mathbf{k}_1} - E_{\mathbf{k}_2} - \epsilon_{\mathbf{k}_1 + \mathbf{k}_2}$.

QUASIPARTICLE RESIDUE

Assuming that the variational wavefunction in Eq. (2) is normalized, $\langle \psi | \psi \rangle = 1$, the residue is given by $Z = |\alpha_0|^2$. Carrying out the renormalization procedure, we arrive at

$$Z = \frac{\left(\frac{\sqrt{n_0}\eta}{E} - \sum_{\mathbf{k}} \frac{v_{\mathbf{k}}\xi_{\mathbf{k}}}{E} \right)^2}{\left(\frac{\sqrt{n_0}\eta}{E} - \sum_{\mathbf{k}} \frac{v_{\mathbf{k}}\xi_{\mathbf{k}}}{E} \right)^2 + \frac{\eta^2}{g^2} + \sum_{\mathbf{k}} \left(\frac{\sqrt{n_0}\xi_{\mathbf{k}} + u_{\mathbf{k}}\eta}{E - \epsilon_{\mathbf{k}}^{\text{im}} - E_{\mathbf{k}}} \right)^2 + \sum_{\mathbf{k}} \frac{\xi_{\mathbf{k}}^2}{g^2} + \sum_{\mathbf{k}_1 \mathbf{k}_2} \frac{(u_{\mathbf{k}_1}\xi_{\mathbf{k}_2} + u_{\mathbf{k}_2}\xi_{\mathbf{k}_1})^2}{2E_{\mathbf{k}_1 \mathbf{k}_2}^2}}. \quad (\text{S3})$$

DECAY RATE

In the decay rate, we need to evaluate $\sum_{\mathbf{k}} |\gamma_{\mathbf{k}}|^2$. This is evaluated similarly to Z above:

$$\sum_{\mathbf{k}} |\gamma_{\mathbf{k}}|^2 = \frac{\sum_{\mathbf{k}} \frac{\xi_{\mathbf{k}}^2}{g^2}}{\left(\frac{\sqrt{n_0}\eta}{E} - \sum_{\mathbf{k}} \frac{v_{\mathbf{k}}\xi_{\mathbf{k}}}{E} \right)^2 + \frac{\eta^2}{g^2} + \sum_{\mathbf{k}} \left(\frac{\sqrt{n_0}\xi_{\mathbf{k}} + u_{\mathbf{k}}\eta}{E - \epsilon_{\mathbf{k}}^{\text{im}} - E_{\mathbf{k}}} \right)^2 + \sum_{\mathbf{k}} \frac{\xi_{\mathbf{k}}^2}{g^2} + \sum_{\mathbf{k}_1 \mathbf{k}_2} \frac{(u_{\mathbf{k}_1}\xi_{\mathbf{k}_2} + u_{\mathbf{k}_2}\xi_{\mathbf{k}_1})^2}{2E_{\mathbf{k}_1 \mathbf{k}_2}^2}}. \quad (\text{S4})$$